

# Dental DANB-CP

DANB's Coronal Polish (CP)

For More Information – Visit link below:

<https://www.examsempire.com/>

**Product Version**

1. Up to Date products, reliable and verified.
2. Questions and Answers in PDF Format.



<https://examsempire.com/>

Visit us at: <https://www.examsempire.com/danb-cp>

# Latest Version: 6.0

## Question: 1

The OSHA Hazard Communication Standard requires orthodontic clinics to update their exposure control plans:

- A. Every two years.
- B. Biannually.
- C. Every three years.
- D. Annually.

**Answer: D**

Explanation:

The correct frequency for updating exposure control plans in orthodontic clinics, according to the OSHA Hazard Communication Standard, is annually. This requirement is part of a broader approach to ensure workplace safety and health, particularly regarding the handling of hazardous materials.

OSHA, the Occupational Safety and Health Administration, mandates that all employers, including those in healthcare settings like orthodontic clinics, regularly review and update their exposure control plans. These plans are critical for outlining procedures and measures designed to protect employees from hazards associated with their work environment, especially those involving potential exposure to bloodborne pathogens and chemical hazards.

The annual review and update of the exposure control plan ensure that the strategies and practices remain effective and relevant. This includes evaluating the effectiveness of existing controls, incorporating new or revised employee positions, updating the list of tasks and procedures, and integrating advancements in technology and safety equipment. For example, in an orthodontic clinic, changes might include updates due to new dental materials or equipment that could pose different health risks compared to those previously evaluated.

Furthermore, keeping the exposure control plan readily accessible is an essential part of compliance. This accessibility ensures that all employees can review the procedures and protections in place, thereby enhancing their understanding and implementation of safety measures. It also facilitates inspections and reviews by OSHA personnel, who can assess compliance and the effectiveness of the plan during their visits.

In summary, the annual updating of exposure control plans as required by OSHA for orthodontic clinics is a vital part of maintaining a safe working environment. This process helps in managing the evolving nature of workplace hazards and ensures that both the health of employees and the operational standards of the clinic adhere to the regulations set forth by health and safety authorities.

## Question: 2

Which of the following is NOT used for detecting irregularities?

- A. Mirror.

- B. Explorer.
- C. Periodontal probe.
- D. Curette.

**Answer: D**

Explanation:

The question posed is asking which of the listed tools is not used for detecting irregularities. The options provided are Mirror, Curette, Explorer, Periodontal probe, and Curette is mentioned multiple times.

A dental mirror is used to visualize parts of the mouth that are difficult to see directly. It allows dentists to examine teeth and gums for signs of decay, gum disease, or other dental issues, thus aiding in the detection of irregularities.

An explorer, often a sharp-pointed tool, is used by dentists to explore the pockets between teeth, around the gum line, and other hard-to-reach areas. It is particularly useful for detecting cavities, cracks, or other abnormalities in the tooth enamel.

A periodontal probe is a thin, rod-like instrument used to measure the depth of periodontal pockets and assess the health of the periodontal tissues. This tool is crucial for detecting periodontal disease and other issues related to the gums.

On the other hand, a curette is primarily used in the process of curettage, which involves the removal of tartar, plaque, and diseased tissue from the teeth and gum pockets. While it is an essential tool in dental cleaning and surgical procedures, it is not specifically designed for the detection of irregularities but rather for the removal of unwanted materials.

Therefore, the correct answer to the question is "Curette," as it is not primarily used for detecting irregularities but for a different purpose in dental treatment. The other tools mentioned are indeed used for detection purposes, making them incorrect choices for this particular question.

### Question: 3

Which of the following statements is true?

- A. Limit fluoride exposure in children where there is sufficient fluoride in the water.
- B. Always limit fluoride exposure in children because of sensitivity.
- C. Children cannot be exposed to enough fluoride to protect their teeth.
- D. Fluoride is not important for children.

**Answer: A**

Explanation:

The question at hand addresses the importance of monitoring fluoride exposure in children, particularly when there is already sufficient fluoride present in their drinking water. The correct answer is: "Limit fluoride exposure in children where there is sufficient fluoride in the water." This statement is true because excessive fluoride intake can lead to dental fluorosis.

Dental fluorosis is a cosmetic condition that affects the teeth during their development before they break through the gums. It is characterized by the appearance of faint white lines or streaks on the teeth and occurs when children consume too much fluoride over long periods while their teeth are developing

under the gums. While mild dental fluorosis can be barely noticeable, more severe forms can cause considerable enamel discoloration and surface irregularities.

Fluoride is indeed important for dental health as it helps in the prevention of cavities by making the tooth enamel more resistant to acid attacks from bacteria in the mouth. However, the benefits of fluoride are dose-dependent, and too much can cause harm, underscoring the importance of balancing fluoride intake.

It is essential for guardians and healthcare providers to ensure that children are not exposed to excessive fluoride, especially in areas where the drinking water is fluoridated. Monitoring other sources of fluoride like toothpaste and dietary supplements is also crucial. Children under the age of six are particularly at risk for fluorosis because their permanent teeth are still forming.

Therefore, while fluoride is critical for dental health, its exposure must be carefully managed to prevent the risk of fluorosis, ensuring children reap the benefits of fluoride without facing the negative effects of overexposure.

### Question: 4

When beginning a cup polishing procedure, which of the following is true?

- A. Begin with the front center teeth and move towards the back of the mouth.
- B. Begin with the distal surface of the most posterior tooth and work towards the front of the mouth.
- C. Begin at the crown and work toward the gingiva.
- D. Begin with the handset on the highest speed.

**Answer: B**

Explanation:

When initiating a cup polishing procedure in dental practice, it is essential to start with the distal surface of the most posterior tooth in the patient's mouth. This strategic approach ensures a systematic and comprehensive cleaning process. By beginning at the distal surface—the surface of the tooth furthest from the midline of the face—you ensure that all surfaces of each tooth are attended to meticulously. The reason for starting at the back of the mouth and working towards the front is to maintain a clear field of view and to systematically remove plaque and debris without redistributing it to already cleaned areas. This method also helps in maintaining the orientation and ensuring that no tooth is missed during the polishing process.

Additionally, it is recommended to begin the procedure using the slowest speed setting on the handpiece. Starting slow allows the dental professional to control the polisher more effectively and minimizes the risk of damaging the tooth enamel or the gingiva. As you become more accustomed to the patient's mouth and the placement of teeth, you can gradually increase the speed of the handpiece to efficiently polish each tooth surface.

By following these guidelines—starting from the most posterior tooth's distal surface and moving anteriorly, and beginning with a slow speed before increasing—you can ensure a thorough and safe cup polishing procedure that enhances oral hygiene without compromising tooth integrity. This methodical approach not only improves the efficacy of the cleaning process but also enhances patient comfort during the procedure.

### Question: 5

Which bony structure has the oral cavity and protects the brain?

- A. Cranium.
- B. Scapula.
- C. Vertebral column.
- D. Rib cage.

**Answer: A**

Explanation:

The correct answer to the question is the cranium. The cranium, also known as the skull, is a complex bony structure that primarily serves to protect the brain from injury. It forms the head's upper and back part and houses the brain, ensuring its safety from external shocks or impacts through its hard, robust structure.

Additionally, the cranium also supports the structures of the face and forms a part of the nasal cavity and orbits (eye sockets), playing a crucial role in the sensory functions of sight, smell, and taste. One of the critical features relevant to this question is that the lower part of the cranium includes the oral cavity or mouth. This is where the teeth are set, and it is an essential aspect for the field of dentistry and orthodontics.

Other options mentioned like the scapula, vertebral column, and rib cage, while important in the skeletal system, do not fulfill the functions related to protecting the brain or housing the oral cavity. The scapula, or shoulder blade, is involved in arm movement and is located at the back of the ribcage. The vertebral column, or spine, supports the body and houses the spinal cord but is not involved with the brain or oral cavity. The rib cage protects the thoracic cavity and supports the respiratory system but again does not relate to brain protection or the oral cavity.

In conclusion, the cranium is the only structure among the options given that encompasses both the protection of the brain and the inclusion of the oral cavity, making it the correct answer to the question.

### Question: 6

Which position may a right-handed take when cleaning the lingual aspect of the mandibular anterior teeth?

- A. 5 o'clock.
- B. 12 o'clock.
- C. 10 o'clock.
- D. 6 o'clock.

**Answer: B**

Explanation:

When discussing the optimal position for a right-handed dental practitioner or dental assistant when cleaning the lingual aspect of the mandibular anterior teeth, the 12 o'clock position is generally considered most effective. This determination of positioning is based on the need for efficiency, visibility, and ergonomics during dental procedures.

The 12 o'clock position refers to the practitioner standing directly behind the patient, who is typically reclined in the dental chair. From this vantage point, the practitioner has a direct and unobstructed view of the mandibular anterior teeth, which includes the lower front teeth. This position allows the practitioner to use their dominant hand (right hand, in this case) effectively, providing the stability and precision needed for dental scaling or cleaning procedures on these specific teeth.

In dentistry, the clock positions are used as a guideline to describe where the practitioner is positioned relative to the patient's head. For right-handed practitioners working on the mandibular anterior teeth, other positions like 8 o'clock or 9 o'clock position them to the side of the patient. These side positions might be used for other dental tasks or for accessing different areas of the mouth, but they do not offer the same direct access and visibility as the 12 o'clock position for accessing the lingual surfaces of the lower front teeth.

The choice between these positions can depend on various factors including the specific dental procedure being performed, the practitioner's comfort and experience, and the patient's comfort. However, for tasks requiring detailed visibility and access to the lingual aspect of the mandibular anterior teeth, the 12 o'clock position is preferable. This position minimizes the physical strain on the practitioner, reducing the risk of ergonomic injuries, and also helps in maintaining a clear view of the working area, which is crucial for effective and safe dental cleaning.

Therefore, when a right-handed practitioner is cleaning the lingual aspect of the mandibular anterior teeth, the 12 o'clock position is the correct and most efficient choice. This positioning ensures optimal control of dental instruments and the best possible patient care.

## Question: 7

Abrasive agents are contraindicated if a patient has \_\_\_\_\_.

- A. Sensitive teeth.
- B. Extrinsic stains.
- C. Plaque.
- D. Tobacco stains.

**Answer: A**

Explanation:

Abrasive agents are contraindicated if a patient has sensitive teeth. This contraindication is due to several reasons related to the condition of the teeth and how abrasive agents work.

Sensitive teeth typically suggest that the protective layer of enamel on the teeth has worn down, exposing the dentin underneath. Dentin contains microscopic tubules that lead directly to the nerve endings. When dentin is exposed, it can result in increased sensitivity to temperature, touch, and acidic or sugary substances.

Abrasive agents, commonly found in various types of toothpaste and dental cleaning treatments, are designed to remove surface stains and plaque through a mechanical scrubbing action. These abrasives, which can include substances like calcium carbonate, hydrated silica, and dicalcium phosphate, are effective in cleaning the outer surface of the teeth. However, they can also contribute to the wear and tear of the enamel especially if it is already thin or compromised, as is the case with sensitive teeth.

Further erosion of the enamel by abrasive agents can expose more of the dentin, thus increasing tooth sensitivity and potentially leading to other dental issues such as cavities or severe tooth decay.

Therefore, in cases of sensitive teeth, dental professionals often recommend using non-abrasive

toothpaste specifically formulated for sensitive teeth. These toothpastes typically contain compounds like potassium nitrate or strontium chloride, which help to block the pathways through the dentin to the nerves, reducing sensitivity rather than removing physical layers of the tooth surface.

It is important for individuals with sensitive teeth to consult with dental professionals before choosing dental care products. These professionals can provide guidance on appropriate products that minimize discomfort and protect against further dental erosion. For cases involving stains like extrinsic stains or tobacco stains, alternative dental treatments that do not involve abrasive agents can be considered, such as professional dental cleaning, bleaching, or the use of enzymatic cleaners that break down stains without mechanical scrubbing.

### Question: 8

What is a sign of poor polishing technique?

- A. Warm teeth.
- B. Gingival lacerations.
- C. Gingival margin injured.
- D. Calculus removal.

**Answer: C**

Explanation:

A sign of poor polishing technique is an injured gingival margin. This occurs when the tools used for polishing teeth are not handled correctly, potentially leading to damage at the base of the tooth where it meets the gumline. The gingival margin is particularly sensitive and susceptible to injury if the dental professional is not careful.

During dental treatments that involve polishing, it is crucial to use a gentle touch and the correct equipment to avoid harming this delicate area of the gums. If the gingival margin is damaged, it can lead to discomfort and increased sensitivity for the patient, potentially complicating the healing process. Moreover, injuries to the gingiva can increase the risk of infection, further exacerbating any dental issues.

Proper technique involves not only using the right tools but also being aware of the pressure applied during polishing. Training and experience play a significant role in executing these procedures safely. Dental professionals must ensure they are continually refining their skills to prevent such injuries.

In summary, an injured gingival margin is a clear indicator of poor polishing technique. It underscores the importance of careful, skilled handling of dental instruments to protect patients' oral health and ensure their comfort during and after the procedure.

### Question: 9

Plaque that exists below the gumline is known as which of the following?

- A. Supragingival.
- B. Superingival.
- C. Subgingival.
- D. Undergingival.

**Answer: C**

Explanation:

Subgingival plaque refers to the dental plaque that accumulates below the gumline, specifically in the area between the gums and the teeth, known as the sulcus or periodontal pockets. This type of plaque is particularly concerning because it is more difficult to remove and can contribute significantly to periodontal disease, including gingivitis and periodontitis.

Contrastingly, supragingival plaque forms above the gumline on the visible parts of the teeth. This plaque is easier to clean through routine oral hygiene practices such as brushing and flossing. The presence of supragingival plaque, however, can lead to the development of subgingival plaque if not properly managed. Bacteria and food particles from supragingival plaque can migrate below the gumline, contributing to the formation and growth of subgingival plaque.

The distinction between subgingival and supragingival plaque is crucial because each type affects oral health differently and requires different approaches for effective management. Subgingival plaque, being below the gumline, often necessitates professional cleaning procedures, such as scaling and root planing, to ensure it is thoroughly removed. These procedures help to manage the plaque and tartar buildup that contribute to gum disease.

Understanding these differences highlights the importance of comprehensive oral hygiene that addresses both supragingival and subgingival plaque to maintain overall dental health and prevent periodontal diseases. Regular dental check-ups and cleanings play a vital role in managing these forms of plaque and maintaining gum health.

### Question: 10

When polishing the lingual aspect of the maxillary left posterior quadrant, which position should a right-handed operator be in?

- A. 10 o'clock.
- B. 8 o'clock.
- C. 9 o'clock.
- D. 2 o'clock.

**Answer: C**

Explanation:

Proper operator positioning is crucial in dental procedures not only for the effectiveness of the treatment but also to ensure the comfort and safety of both the patient and the operator. In dental terminology, operator positions around the patient are often described in terms of a clock face for standardization and ease of understanding.

When discussing the right-handed operator working on the lingual aspect (the side of the teeth closest to the tongue) of the maxillary left posterior quadrant (which includes teeth like the molars and premolars on the upper left side of the patient's mouth), positioning is key to accessing the area efficiently and comfortably. For a right-handed operator, the 9 o'clock position is recommended. This position means that the operator is essentially sitting or standing facing the left side of the patient's head, with the patient's nose pointing towards 12 o'clock.

At the 9 o'clock position, the right-handed operator has optimal ergonomic access to the target area. This position allows the operator's dominant hand to access the lingual surfaces of the upper left teeth without strain. The operator's hand and wrist can maintain a natural alignment, reducing the risk of fatigue and repetitive strain injuries. Furthermore, visibility is maximized as the operator can look directly into the quadrant without having to twist awkwardly or use mirrors excessively.

This position contrasts with other potential positions: - At 10 o'clock, while still feasible, the angle becomes slightly more awkward for accessing the left side as it would slightly favor the operator's non-dominant hand. - At 8 o'clock, the operator would be too far behind the patient, making it difficult to reach the lingual surfaces effectively. - At 2 o'clock, which is ideal for working on the right side of the patient's mouth, would be highly impractical for the left side as it would require the operator to reach over the patient's face, causing discomfort and compromising the control over the dental instruments. Hence, the 9 o'clock position is strategically the best choice for a right-handed operator working on the lingual aspect of the maxillary left posterior quadrant. This position ensures optimal access, visibility, ergonomic efficiency, and patient comfort during the dental procedure.

**Thank You for Trying Our Product**  
**Special 16 USD Discount Coupon: NSZUBG3X**

**Email:** [support@examsempire.com](mailto:support@examsempire.com)

**Check our Customer Testimonials and ratings  
available on every product page.**

**Visit our website.**

**<https://examsempire.com/>**