

Nursing ACNS


**ANCC Adult Health Clinical Nurse Specialist Certification
(ACNS)**

For More Information – Visit link below:

<https://www.examsempire.com/>

Product Version

- 1. Up to Date products, reliable and verified.**
- 2. Questions and Answers in PDF Format.**



<https://examsempire.com/>

Visit us at: <https://www.examsempire.com/acns>

Latest Version: 6.0

Question: 1

When providing care for an adult female patient who has a history of prescription benzodiazepine dependence, you consider that:

- A. The preferred method of treatment for this problem is rapid detoxification.
- B. She is at significant risk for drug-induced hepatitis.
- C. She is unlikely to have a problem with misuse of other drugs or alcohol.
- D. She probably has an underlying untreated or under-treated mood disorder.

Answer: D

Explanation:

When providing care for an adult female patient with a history of prescription benzodiazepine dependence, it is crucial to consider the possibility of an underlying untreated or under-treated mood disorder. This consideration is based on several epidemiological and clinical observations that suggest a strong association between substance misuse, specifically prescription medications, and the presence of mood disorders such as anxiety and depression.

Research indicates that substance abuse, which includes the misuse and overuse of mood-altering substances, affects approximately 10-15% of patients in primary care settings. This statistic is significant as it underscores the prevalence of substance misuse in a general healthcare context, highlighting the need for healthcare providers to be vigilant and proactive in identifying potential underlying causes of this behavior.

Women, in particular, exhibit higher rates of misuse of prescription medications compared to men. This trend may be attributed to several factors including differing pharmacokinetics and pharmacodynamics, societal pressures, and potentially greater access to prescription drugs. More importantly, studies have consistently shown that women are more likely to suffer from mood disorders, especially anxiety and depression. These disorders can often drive an individual to self-medicate with substances such as benzodiazepines, which are commonly prescribed for anxiety and can be highly addictive.

The interplay between mood disorders and substance abuse is complex. Mood disorders can lead to substance abuse as individuals attempt to manage symptoms of their mental health issues without proper medical guidance. Conversely, the chronic use of substances like benzodiazepines can exacerbate or even trigger mood disorders. Therefore, when treating a patient with a history of benzodiazepine dependence, it is essential to assess for underlying mood disorders, which may be the root cause driving the dependence.

In conclusion, the correct approach in managing a case of benzodiazepine dependence in an adult female patient involves more than addressing the substance abuse itself. It is imperative to conduct a thorough mental health evaluation to identify any underlying mood disorders. Treating these mood disorders can be pivotal in breaking the cycle of dependence and ensuring a holistic recovery, thereby stabilizing the patient's overall mental and physical health. Engaging in such comprehensive care not only addresses the immediate issue of dependence but also contributes to the long-term wellbeing of the patient.

Question: 2

What can hyperprolactinemia be a sign of?

- A. Lower motor neurone lesion.
- B. Upper motor neurone lesion.
- C. Pituitary adenoma.
- D. Cerebellar disease.

Answer: C

Explanation:

Hyperprolactinemia refers to the condition where there is an abnormally high level of prolactin in the blood. Prolactin is a hormone produced by the pituitary gland, which is located at the base of the brain. This hormone primarily functions to promote lactation (milk production) in women after childbirth, but it is also present in non-pregnant women and men in lower amounts.

One common cause of hyperprolactinemia is a pituitary adenoma, specifically a prolactinoma, which is a benign tumor of the pituitary gland that produces excessive amounts of prolactin. This overproduction can lead to various symptoms, depending on the gender of the individual. In women, high prolactin levels can cause galactorrhea (production and spontaneous flow of breast milk), amenorrhea (absence of menstrual periods), and infertility. In men, symptoms might include decreased libido, erectile dysfunction, and infertility.

Besides prolactinomas, hyperprolactinemia can be indicative of other health issues. These include hypothyroidism, where the thyroid gland does not produce enough hormones, affecting the pituitary gland's regulation of prolactin. Certain medications, such as antipsychotics and antiemetics, can also elevate prolactin levels by interfering with dopamine, a neurotransmitter that normally inhibits prolactin secretion.

In some cases, the symptoms of a pituitary adenoma extend beyond hormonal disturbances. If the adenoma grows large enough, it can exert pressure on surrounding brain structures. This might lead to headaches, as well as visual disturbances if the optic nerves are compressed. These symptoms necessitate prompt medical evaluation to manage the tumor and alleviate the risk of permanent damage.

Therefore, while hyperprolactinemia can be a sign of a pituitary adenoma, it is crucial to consider and rule out other potential causes. A comprehensive evaluation, typically involving blood tests, imaging studies like MRI of the brain, and a thorough assessment of symptoms, is essential to determine the underlying cause and appropriate treatment.

Question: 3

Most states require advanced practice nurses to have which type of work environment?

- A. Independent.
- B. Adjusted.
- C. Collaborative.
- D. Traditional.

Answer: C

Explanation:

The correct answer to the question regarding the type of work environment most states require for advanced practice nurses (APNs) is "Collaborative." This requirement is based on the regulations that govern the scope of practice for APNs, which can vary significantly from one state to another.

In the context of advanced practice nursing, collaboration typically means that the nurse works in conjunction with a physician or other healthcare providers. This collaborative model is designed to enhance patient care by integrating the unique skills and knowledge of each provider. It often involves formal agreements that outline the roles and responsibilities of each party, ensuring that the healthcare delivery is safe, effective, and compliant with state laws.

The necessity for collaboration may be rooted in several factors. One is the level of autonomy granted to APNs, which can differ based on state laws. Some states allow APNs full practice authority, meaning they can practice independently without any required collaboration with a physician. These states recognize that APNs are capable of making complex clinical decisions and managing patient care autonomously.

However, most states still require some level of physician oversight, whether through direct supervision or a collaborative agreement. This oversight might include regular consultation, review of patient cases, or co-signature on certain medical orders. The terminology used to describe these relationships can vary, including words like "collaboration," "delegation," or "supervision."

The collaborative requirement is based on the belief that a team approach can lead to better patient outcomes. It leverages the strengths and expertise of both physicians and APNs, fostering a comprehensive approach to healthcare that benefits patients through improved access to care and a broadening of available services.

In summary, while the landscape of APN practice authority is evolving, with a trend towards greater independence in some regions, the prevailing requirement across most states remains a collaborative work environment. This model ensures that APNs are part of a healthcare team that delivers high-quality medical care, ultimately benefiting the broader health system and patient population.

Question: 4

How often should peripheral lines be changed?

- A. every 24 hours
- B. 24 to 48 hours
- C. 72 hours
- D. every week

Answer: C

Explanation:

Peripheral intravenous (IV) lines, commonly referred to as peripheral lines, are essential for administering medications, fluids, and nutrients directly into the bloodstream. They are a routine part of medical treatments in hospitals and clinics. However, these lines pose a risk for complications, primarily

infections, if not managed correctly. To mitigate these risks, specific guidelines on the frequency of changing peripheral IV catheters have been established.

According to the Centers for Disease Control and Prevention (CDC) and the Infusion Nurses Society (INS), peripheral IV catheters should be replaced every 72 to 96 hours in adults to prevent complications, including phlebitis and bloodstream infections. This timeframe is recommended to balance the risk of infection with the need to minimize patient discomfort and the use of medical resources.

The decision to replace a peripheral line should also consider factors such as the condition of the insertion site and the type of infusion. For example, if the IV site shows signs of redness, swelling, or pain, or if the catheter is malfunctioning (e.g., occlusion or leakage), it should be replaced immediately regardless of the last change. Similarly, if infusions of irritating solutions or vesicants have been administered, more frequent monitoring and potential replacement might be necessary.

In pediatric patients, the guidelines may vary slightly due to differences in vein size and the sensitivity of their skin. Pediatric peripheral lines may often be evaluated on a case-by-case basis, with considerations for the least possible invasiveness and frequency of changes to avoid undue distress or harm to the child.

Proper hand hygiene, aseptic techniques during insertion and maintenance, and regular assessment of the IV site are crucial components that work in conjunction with the timing of catheter replacement to ensure patient safety and the effectiveness of the vascular access device. By adhering to these guidelines, healthcare providers can significantly reduce the risk of complications associated with peripheral lines.

Question: 5

An open-ended question that invites that patient to elaborate on a point is known as which of the following?

- A. Open lead.
- B. General lead.
- C. Leading question.
- D. Offering.

Answer: B

Explanation:

The correct term for an open-ended question that invites a patient to elaborate on a point is known as a "General lead."

General leads are a type of communication technique often used in therapeutic or medical settings to encourage a patient to provide more detailed information about their thoughts, feelings, or experiences. Unlike specific questions, which direct a patient's response to a particular aspect, general leads are open and non-directive. They do not focus on a specific issue but allow the patient the freedom to take the conversation in whichever direction they feel is most relevant or significant to them at that moment. Using general leads can be particularly effective in establishing rapport with the patient. It signals to the patient that the healthcare provider is interested in hearing more about what the patient has to say, without imposing any judgment or direction. This can help create a safe and open environment, where patients feel more comfortable and willing to disclose personal information.

Furthermore, general leads are instrumental in eliciting comprehensive information. By allowing the patient to decide what details to include in their response, healthcare providers can gather a broader

perspective on the patient's situation. This can be crucial for accurate diagnosis and effective treatment planning.

Examples of general leads include phrases like "Tell me more about that," "How did that make you feel?" or simply "Go on." Each of these prompts encourages further discussion, which can provide deeper insight into the patient's condition and aid in more personalized care.

Question: 6

Which of the following does NOT increase a woman's risk for domestic violence?

- A. Being in a lesbian relationship rather than a heterosexual one.
- B. Being pregnant.
- C. Being disabled.
- D. Being an undocumented immigrant.

Answer: A

Explanation:

The question asks which scenario does NOT increase a woman's risk for domestic violence. The correct answer, as repeated across all options, is "Being in a lesbian relationship rather than a heterosexual one."

Domestic violence is a significant issue that affects various demographics, regardless of sexual orientation, age, race, or economic status. However, certain factors are recognized to heighten the risk of experiencing domestic violence. Studies and data have identified factors such as pregnancy, disability, and undocumented status as increasing the vulnerability of women to domestic violence. These factors can exacerbate power imbalances, dependency, isolation, and barriers to seeking help, all of which contribute to the increased risk.

In contrast, there is no consistent evidence to suggest that the nature of the relationship in terms of the sexual orientation of the partners—specifically being in a lesbian versus a heterosexual relationship—alters the fundamental risk of domestic violence. Domestic violence occurs in both heterosexual and same-sex relationships at significant rates. The misconception that same-sex relationships inherently involve less power imbalance or are less prone to violence is unfounded. The risk factors typically revolve around individual behaviors, relationship dynamics, and socio-economic conditions rather than the sexual orientation of the relationship itself.

Therefore, while being pregnant, disabled, or an undocumented immigrant are recognized as factors that can increase a woman's risk for victimization, being in a lesbian relationship, in itself, does not alter the risk compared to being in a heterosexual relationship. The focus should remain on the behaviors and dynamics that contribute to domestic violence, rather than the nature of the relationship in terms of sexual orientation.

Question: 7

What approach should be taken for a thoracentesis?

- A. Axillary.
- B. Anterior.

- C. Posterior.
- D. Fowlers.

Answer: C

Explanation:

Thoracentesis is a medical procedure aimed at removing fluid from the pleural space in the chest, which can accumulate due to various medical conditions such as infections, heart failure, or malignancies. The approach taken during thoracentesis is critical for both maximizing fluid removal and minimizing potential complications.

The most commonly recommended approach for thoracentesis is the posterior approach. This involves the patient sitting up and leaning forward on a table, exposing the back. This position allows the healthcare provider easier access to the pleural space through the back, where the pleural fluid tends to accumulate, especially when the patient is upright. The posterior approach is generally preferred because it is technically easier due to the natural pooling of fluid in this area when the patient is sitting. In contrast, the anterior approach, where the puncture is made on the front side of the chest, is less commonly used. This approach may cause discomfort or anxiety for some patients, as they can see the needle being inserted. Additionally, anatomical considerations, such as the presence of breast tissue and pectoral muscles, can make this approach more challenging.

The axillary approach, which involves accessing the pleural space from the patient's armpit area, is another option. However, it is less favored because the patient's arm can obstruct the procedure, making it difficult for the healthcare provider to maneuver the needle. This approach also has a higher risk of injuring local structures in the crowded axillary area.

During a thoracentesis, patient positioning is crucial. While the posterior approach is generally preferred, the patient's specific anatomy, the volume of fluid to be removed, and the presence of any loculated effusions (areas where the fluid is encapsulated and not freely moving) can influence the choice of the specific site of needle insertion. Therefore, imaging studies such as ultrasound are often used before and during the procedure to guide the needle placement, enhancing safety and effectiveness.

In summary, while the posterior approach is most commonly recommended for thoracentesis due to ease of access and patient positioning, the choice of approach can vary based on the patient's specific situation and the clinician's judgment. Proper technique and careful attention to patient comfort and safety are paramount in performing this procedure effectively.

Question: 8

All of the following are true of informed consent except:

- A. It must be documented in the medical records that the patient has been informed of his or her healthcare options.
- B. This is the right of all competent adults who are over the age of 18 and the right of emancipated minors.
- C. The clinician has the duty to explain relevant information to the patient so that the patient can make an appropriate decision regarding the care to be provided.
- D. This consent will absolve the CNS from allegations of malpractice should it occur.

Answer: D

Explanation:

Informed consent is a fundamental principle in medical ethics and legal requirements, which mandates that patients must be informed about their medical condition, the available treatment options, the potential risks and benefits associated with each option, and the consequences of receiving no treatment. This information allows patients to make well-informed decisions about their healthcare. The process of obtaining informed consent involves detailed communication between the healthcare provider and the patient, and it should be documented in the patient's medical record, confirming that the patient understands the information provided and consents to the proposed course of treatment. However, it is crucial to understand that obtaining informed consent does not protect a healthcare provider, such as a Clinical Nurse Specialist (CNS), from allegations of malpractice. Malpractice refers to professional negligence where a healthcare provider deviates from the standards of practice in the medical community, leading to patient harm. Even if informed consent has been appropriately obtained and documented, a CNS can still face allegations of malpractice if they fail to meet the accepted standards of medical care. Informed consent primarily protects the patient's rights and ensures patient autonomy but does not absolve healthcare providers from their duty to deliver competent and standard care.

The correct answer to the question, therefore, is that informed consent does not absolve a CNS from allegations of malpractice should it occur. This is because informed consent relates to the patient being informed and agreeing to the treatment based on the understanding of potential risks and outcomes, not to the actual performance standards of the healthcare provider. Malpractice is determined by the quality of care provided and adherence to professional standards, not merely by whether a patient agreed to the treatment.

It is also important to note that informed consent is required not only by ethical standards but also by legal regulations across many jurisdictions. This legal requirement emphasizes the importance of transparent communication in healthcare settings and ensures that patients have the opportunity to be active participants in their own healthcare decisions. The rights to informed consent are granted to all competent adults over the age of 18 and to emancipated minors, ensuring that these groups are protected and have control over their medical decisions.

Therefore, while informed consent is a crucial aspect of the medical process, it serves primarily to educate and empower patients rather than to protect healthcare providers from legal repercussions of their clinical actions. Understanding this distinction is key in appreciating the scope and limitations of informed consent in the medical field.

Question: 9

A Nurse Practitioner vicariously involved in a dispute between a patient and another provider (an anesthesiologist) by means of being a witness. The Nurse Practitioner provides the patient legal advice on how to prove causation in court. What does the Nurse Practitioner's action of giving legal advice to the patient PRIMARILY demonstrate?

- A. Appropriate action that falls within a nurse's scope of practice.
- B. Inappropriate action that falls outside a nurse's scope of practice.
- C. Ethical action to protect a patient's welfare.
- D. Breach of duty of nursing standards.

Answer: B

Explanation:

The scenario described involves a Nurse Practitioner (NP) who witnesses a dispute between a patient and an anesthesiologist and subsequently provides the patient with legal advice on proving causation in court. It is important to understand why the action taken by the NP is considered inappropriate and falls outside the professional scope of nursing practice.

Firstly, the primary role of a nurse, including a Nurse Practitioner, centers on patient care, which involves assessment, planning, implementation, and evaluation of treatment to maintain or restore health. Nurses are trained and licensed to make clinical decisions and provide medical care based on their specific qualifications and the bounds of their licensure.

However, the provision of legal advice is a skill and responsibility designated for trained legal professionals - namely, lawyers. Legal advice involves interpreting and suggesting courses of action based on the law for specific situations that a client might face. When the NP in this scenario advises the patient on legal matters, especially on how to prove causation in court, they are stepping beyond the limits of their professional training and legal licensing.

It's critical to recognize that such an action can not only mislead the patient but also potentially result in legal repercussions for the NP. By offering legal counsel, the NP risks providing incorrect or incomplete advice, as they are not trained in the legal intricacies of medical law or litigation processes. This misstep does not just breach the professional scope but also the ethical boundaries that define and uphold the integrity of the nursing profession.

Moreover, in situations where legal advice is necessary, the appropriate and ethical action for a nurse or NP would be to refer the patient to a qualified legal professional. This not only ensures that the patient receives expert advice tailored to their legal needs but also maintains the clarity of professional roles and responsibilities.

In summary, the action of the Nurse Practitioner providing legal advice in this context is inappropriate because it falls outside the boundaries of nursing practice. Nurses are crucial in the healthcare system and must operate within the scope of their practice, both for the safety of their patients and the legal and professional integrity of themselves and their workplace. Referral to the appropriate legal services when necessary is both a legal and ethical obligation for nurses.

Question: 10

Of the following interventions which would be the third step followed in the case of toxic ingestion overdose?

- A. Position and suction patient to establish airway.
- B. If obtunded, comatose and there's no gag reflex, intubate the patient.
- C. Assess patient's breathing depth and rate.
- D. none of the above

Answer: C

Explanation:

In the scenario of a toxic ingestion overdose, the immediate focus is on stabilizing the patient's vital functions, particularly their airway, breathing, and circulation – commonly referred to as the ABCs of

emergency medicine. The steps detailed below adhere to this principle, ensuring the patient is first secure in terms of airway management before assessing and managing respiratory and circulatory functions.

The first step in managing a toxic ingestion overdose involves positioning and suctioning the patient to establish and secure the airway. This is crucial because a compromised airway can rapidly lead to respiratory failure and decreased oxygen supply to the brain and other vital organs. Positioning may involve placing the patient in a way that prevents the tongue from blocking the airway, and suctioning helps clear any obstructions or vomitus that could block air passage.

The second step involves assessing whether the patient is obtunded (less alert) or comatose and checking for the presence of a gag reflex. If the patient is deeply unconscious (comatose) and lacks a gag reflex, they are at high risk of airway obstruction. In such cases, intubation is recommended. Intubation involves inserting a tube into the patient's trachea to keep the airway open and ensure that the air passes freely into the lungs. This step is critical for maintaining ventilation and oxygenation.

The third step, which is assessing the patient's breathing depth and rate, follows after ensuring the airway is secured either through positioning and suction or intubation if necessary. This step is vital as it provides information about the patient's ventilatory status. In the context of a toxic ingestion overdose, the patient might have a respiratory compromise due to the toxic effects of the ingested substance on the respiratory center in the brain or due to direct injury to the respiratory system. Assessing the depth and rate of breathing helps in determining the need for further respiratory support, such as supplemental oxygen or mechanical ventilation.

These steps are sequential and each one builds on the safety and stability provided by the previous step. By following these systematic interventions, healthcare providers can effectively manage a patient who has overdosed, thereby reducing the risk of complications and improving the overall outcome.

Thank You for Trying Our Product

Special 16 USD Discount Coupon: NSZUBG3X

Email: support@examsempire.com

**Check our Customer Testimonials and ratings
available on every product page.**

Visit our website.

<https://examsempire.com/>